

Chaplygin gas with non-adiabatic pressure perturbations

Winfried Zimdahl^{†‡} and Julio C. Fabris^{‡§}

[†] Institut für Theoretische Physik, Universität zu Köln, D-50937 Köln, Germany

[‡] Departamento de Física Universidade Federal do Espírito Santo,
CEP29060-900 Vitória, Espírito Santo, Brazil

Abstract. Perturbations in a Chaplygin gas, characterized by an equation of state $p = -A/\rho$, may acquire non-adiabatic contributions if spatial variations of the parameter A are admitted. This feature is shown to be related to a specific internal structure of the Chaplygin gas. We investigate how perturbations of this type modify the adiabatic sound speed and influence the time dependence of the gravitational potential which gives rise to the Integrated Sachs-Wolfe effect in the anisotropy spectrum of the cosmic microwave background.

[†] Electronic address: zimdahl@thp.uni-koeln.de

[§] Electronic address: fabris@cce.ufes.br

1. Introduction

Since 1998 [1] a growing number of observational data has backed up the conclusion that the expansion rate of our present Universe is increasing. According to our current understanding on the basis of Einstein's General Relativity, such a dynamics requires a cosmic substratum with an effective negative pressure. To clarify the physical nature of this substratum is one of the major challenges in cosmology. Most of the approaches in the field rely on a two-component picture of the cosmic medium with the (at present) dynamically dominating "Dark Energy" (DE), equipped with a negative pressure, which contributes roughly 70% to the total energy density, and with pressureless (Cold) "Dark Matter" (CDM), which contributes roughly 30% (see, e.g. [2] and references therein). Usually, these components are assumed to evolve independently, but more general interacting models have been considered as well [3]. A single-component model which has attracted some interest as an alternative description is a Chaplygin gas [4]. The Chaplygin gas which is theoretically based in higher dimensional theories [5], has been considered as a candidate for a unified description of dark energy and dark matter [6, 7, 8, 9, 10]. Its energy density smoothly changes from that of matter at early times to an almost constant value at late times. It interpolates between a phase of decelerated expansion, necessary for structure formation to occur, and a subsequent period in which the dynamically dominating substratum acts similarly as a cosmological constant, giving rise to accelerated expansion. The Chaplygin gas combines a negative pressure with a positive sound velocity. This sound speed is negligible at early times and approaches the speed of light in the late-time limit. A sound speed of the order of the speed of light has implications which apparently disfavor the Chaplygin gas as a useful model of the cosmic medium. In particular, it should be connected with oscillations of the medium on small (sub-horizon) scales. The fact that the latter are not observed has led the authors of [11] to the conclusion that Chaplygin gas models of the cosmic medium are ruled out as competitive candidates. Similar results were obtained from the analysis of the anisotropy spectrum of the cosmic microwave background [12, 13], except possibly for low values of the Hubble parameter [14]. However, these conclusions rely on the assumption of an adiabatic cosmic medium. It has been argued that there might exist entropy perturbations, so far not taken into account, which may change the result of the adiabatic perturbation analysis [15, 16]. A problem here is the origin of non-adiabatic perturbations which should reflect the internal structure of the cosmic medium. The latter is unknown but it may well be more complicated than suggested by the usually applied simple (adiabatic) equations of state. Generally, non-adiabatic perturbations will modify the adiabatic sound speed. The speed of sound has attracted interest as a tool to discriminate between different dark energy models [17, 18, 19]. It has been argued that (anti-)correlations between curvature and isocurvature fluctuations may provide a mechanism to modify the (adiabatic) sound speed of the dark energy [20, 22]. A reduced sound speed (compared with the velocity of light) in quintessence models may lead to a suppression of the lowest multipole moments of the CMB anisotropy spectrum

(which are usually enhanced by the Integrated Sachs-Wolfe (ISW) effect [18, 21, 22]). The point here is, that a change of the sound speed will affect the time dependence of the gravitational potential on large perturbation scales. It is this time-dependence that gives rise to the ISW effect [23]. A detection of the ISW, representing an independent confirmation of the existence of dark energy, has recently been reported by [24, 25, 26]. Since the CMB data suggest reduced power of the lowest multipoles (compared with the prediction of the Λ CDM model), different reduction mechanisms have been discussed in the literature (see, e.g. [27]), among them also approaches that require “new physics” (see, e.g., [28, 29]), which will not be considered, however, in this paper.

The purpose of this paper is to study a simple model of non-adiabatic perturbations in a Chaplygin gas. Starting with a two-component description of the cosmic medium we demonstrate that any Chaplygin gas in a homogeneous and isotropic Universe can be regarded as being composed of another Chaplygin gas which is in interaction with a pressureless fluid such that the background ratio of the energy densities of both components is constant. Spatial perturbations of this ratio then give rise to non-adiabatic pressure perturbations which may be related to fluctuations of the equation of state parameter A . We investigate the influence of these perturbations on the sound speed and on the time dependence of the gravitational potential.

The paper is organized as follows. In Section 2 we introduce the necessary definitions and relations to describe the dynamics of an interacting two-fluid model both in a homogeneous and isotropic background and on the perturbative level. Section 3 starts by recalling basic relations for the Chaplygin gas. Then we show that any one-component Chaplygin gas in a homogeneous and isotropic universe is equivalent to an interacting two-component system, where one of the components is another (different parameters) Chaplygin gas and the second component is dust. The interaction is such that the ratio of the energy densities of both components remains constant in time. On the basis of this two-component interpretation we introduce non-adiabatic pressure perturbations in section 4 by allowing the ratio of the energy densities of both components to fluctuate. Relating these fluctuations of the density ratio to fluctuations of the equation of state parameter of the original one-component Chaplygin gas provides us with modifications of the adiabatic sound speed. Furthermore, we investigate the large scale perturbation dynamics and obtain a non-adiabatic contribution to the gravitational potential which is relevant for the ISW. The results of the paper are summarized in Section 5.

2. Interacting two-fluid dynamics

Much of the appeal of a Chaplygin gas model of the cosmic substratum is due to the circumstance that it represents a unified description of dark energy and dark matter within a one component model. In order to describe non-adiabatic features which are the manifestation of an internal structure of the medium it will be instructive, however, to start the discussion by establishing basic relations of a two-fluid description. The point here is that we shall reveal an equivalent two-component picture of any one-

component Chaplygin gas (for different two-component decompositions see [10, 30]). This circumstance can be used to introduce a simple model for an internal structure which gives rise to non-adiabatic pressure perturbations.

2.1. General setting

We assume the cosmic medium to behave as a perfect fluid with an energy-momentum tensor

$$T^{ik} = \rho u^i u^k + p h^{ik}, \quad h^{ik} = g^{ik} + u^i u^k. \quad (1)$$

Our approach is based on the decomposition

$$T^{ik} = T_1^{ik} + T_2^{ik} \quad (2)$$

of the total energy-momentum into two parts with ($A = 1, 2$)

$$T_A^{ik} = \rho_A u_A^i u_A^k + p_A^* h_A^{ik}, \quad h_A^{ik} = g^{ik} + u_A^i u_A^k, \quad T_{A;k}^{ik} = 0. \quad (3)$$

For our purpose it is convenient to split the (effective) pressures p_1^* and p_2^* according to

$$p_1^* \equiv p_1 + \Pi_1, \quad p_2^* \equiv p_2 + \Pi_2. \quad (4)$$

This procedure allows us to introduce an arbitrary coupling between both fluids by the requirement

$$\Pi_1 = -\Pi_2 \equiv \Pi \quad \Leftrightarrow \quad p_1^* = p_1 + \Pi, \quad p_2^* = p_2 - \Pi. \quad (5)$$

Under this condition the apparently separate energy-momentum conservation $T_{A;k}^{ik} = 0$ in (3) is only formal. Then, for a homogeneous and isotropic universe with $u_1^i = u_2^i = u^i$ and $u_{;a}^a = 3H$, where H is the Hubble rate, the individual energy balance equations are coupled and take the form

$$\dot{\rho}_1 + 3H(\rho_1 + p_1) = -3H\Pi \quad \Leftrightarrow \quad \dot{\rho}_1 + 3H(\rho_1 + p_1^*) = 0, \quad (6)$$

and

$$\dot{\rho}_2 + 3H(\rho_2 + p_2) = 3H\Pi \quad \Leftrightarrow \quad \dot{\rho}_2 + 3H(\rho_2 + p_2^*) = 0. \quad (7)$$

Adding up these balances, consistently reproduces the total energy conservation

$$\dot{\rho} + 3H(\rho + p) = 0 \quad (8)$$

with

$$\rho = \rho_1 + \rho_2, \quad p = p_1 + p_2 = p_1^* + p_2^*. \quad (9)$$

The total adiabatic sound velocity may be split according to

$$\frac{\dot{p}}{\dot{\rho}} = \frac{\dot{\rho}_1}{\dot{\rho}} \frac{\dot{p}_1}{\dot{\rho}_1} + \frac{\dot{\rho}_2}{\dot{\rho}} \frac{\dot{p}_2}{\dot{\rho}_2} = \frac{\dot{\rho}_1}{\dot{\rho}} \frac{\dot{p}_1^*}{\dot{\rho}_1} + \frac{\dot{\rho}_2}{\dot{\rho}} \frac{\dot{p}_2^*}{\dot{\rho}_2}. \quad (10)$$

We emphasize that so far the equations of state and the nature of the interaction between the components are left unspecified.

2.2. Matter perturbations

To study perturbations about the homogeneous and isotropic background we introduce the following quantities. We define the total fractional energy density perturbation

$$D \equiv \frac{\hat{\rho}}{\rho + p} = -3H \frac{\hat{\rho}}{\dot{\rho}}, \quad (11)$$

and the energy density perturbations for the components ($A = 1, 2$)

$$D_A \equiv \frac{\hat{\rho}_A}{\rho_A + p_A^*} = -3H \frac{\hat{\rho}_A}{\dot{\rho}_A}. \quad (12)$$

Likewise, the corresponding pressure perturbations are

$$P \equiv \frac{\hat{p}}{\rho + p} = -3H \frac{\hat{p}}{\dot{\rho}} = -3H \frac{\dot{p}}{\dot{\rho}} \frac{\hat{p}}{\dot{p}}, \quad (13)$$

and

$$P_A \equiv \frac{\hat{p}_A}{\rho_A + p_A^*} = -3H \frac{\hat{p}_A}{\dot{\rho}_A} = -3H \frac{\dot{p}_A}{\dot{\rho}_A} \frac{\hat{p}_A}{\dot{p}_A}. \quad (14)$$

One also realizes that

$$D = \frac{\dot{\rho}_1}{\dot{\rho}} D_1 + \frac{\dot{\rho}_2}{\dot{\rho}} D_2 \quad (15)$$

and

$$P = \frac{\dot{\rho}_1}{\dot{\rho}} P_1 + \frac{\dot{\rho}_2}{\dot{\rho}} P_2. \quad (16)$$

In terms of these quantities the non-adiabatic pressure perturbations

$$P - \frac{\dot{p}}{\dot{\rho}} D = -3H \frac{\dot{p}}{\dot{\rho}} \left[\frac{\hat{p}}{\dot{p}} - \frac{\hat{\rho}}{\dot{\rho}} \right] \quad (17)$$

are then generally characterized by

$$\begin{aligned} P - \frac{\dot{p}}{\dot{\rho}} D &= \frac{\dot{\rho}_1}{\dot{\rho}} \left(P_1 - \frac{\dot{p}_1}{\dot{\rho}_1} D_1 \right) + \frac{\dot{\rho}_2}{\dot{\rho}} \left(P_2 - \frac{\dot{p}_2}{\dot{\rho}_2} D_2 \right) \\ &\quad + \frac{\dot{\rho}_1 \dot{\rho}_2}{(\dot{\rho})^2} \left[\frac{\dot{p}_2}{\dot{\rho}_2} - \frac{\dot{p}_1}{\dot{\rho}_1} \right] [D_2 - D_1]. \end{aligned} \quad (18)$$

The first two terms on the right-hand side describe internal non-adiabatic perturbations within the individual components. The last term takes into account non-adiabatic perturbations due to the two-component nature of the medium.

2.3. Metric perturbations

A homogenous and isotropic background universe with scalar metric perturbations is characterized by the line element

$$ds^2 = -(1 + 2\phi) dt^2 + 2a^2 F_{,\alpha} dt dx^\alpha + a^2 [(1 - 2\psi) \delta_{\alpha\beta} + 2E_{,\alpha\beta}] dx^\alpha dx^\beta. \quad (19)$$

The components of the perturbed 4-velocity are

$$\hat{u}^0 = \hat{u}_0 = -\phi \quad (20)$$

and

$$a^2 \hat{u}^\mu + a^2 F_{,\mu} = \hat{u}_\mu \equiv v_{,\mu} . \quad (21)$$

Furthermore, we define

$$\chi \equiv a^2 (\dot{E} - F) . \quad (22)$$

The perturbation dynamics is most conveniently described in terms of the gauge-invariant variable [31]

$$\zeta \equiv -\psi + \frac{1}{3} \frac{\hat{\rho}}{\rho + p} = -\psi - H \frac{\hat{\rho}}{\dot{\rho}} . \quad (23)$$

Corresponding quantities for the components are

$$\zeta_A \equiv -\psi + \frac{1}{3} \frac{\hat{\rho}_A}{\rho_A + p_A^*} = -\psi - H \frac{\hat{\rho}_A}{\dot{\rho}_A} . \quad (24)$$

The variable ζ obeys the equation (cf [32, 33, 34])

$$\dot{\zeta} = -H \left(P - \frac{\dot{p}}{\dot{\rho}} D \right) - \frac{1}{3a^2} \Delta v^x , \quad (25)$$

where

$$v^x \equiv v + \chi \quad (26)$$

is the gauge-invariant velocity potential. Using in (25) the formula (18) for $P - (\dot{p}/\dot{\rho})D$ together with the definitions (24) and taking into account that $p_1 = p_1(\rho_1)$ and $p_2 = p_2(\rho_2)$, Eq. (25) reduces to

$$\dot{\zeta} = -3H \frac{\dot{\rho}_1 \dot{\rho}_2}{\dot{\rho}^2} \left[\frac{\dot{p}_2}{\dot{\rho}_2} - \frac{\dot{p}_1}{\dot{\rho}_1} \right] (\zeta_2 - \zeta_1) - \frac{1}{3a^2} \Delta v^x . \quad (27)$$

Equation (27) expresses the general fact that entropy perturbations $(\zeta_2 - \zeta_1)$ source curvature perturbations ζ . This completes our preliminary definitions and remarks which, under the given conditions, are generally valid. No specific Chaplygin gas properties have been used so far.

3. The Chaplygin gas

3.1. Two-component interpretation

The Chaplygin gas is characterized by an equation of state

$$p = -\frac{A}{\rho} , \quad (28)$$

where p is the pressure and ρ is the energy density of the gas. A is a positive constant. This equation of state gives rise to the energy density [6]

$$\rho = \sqrt{A + \frac{B}{a^6}} , \quad (29)$$

where B is another (positive) constant. Now, let us rename the constants according to

$$A = A_2 (1 + \kappa) , \quad B = B_2 (1 + \kappa)^2 . \quad (30)$$

For any constant, non-negative κ the quantities A_2 and B_2 are constant and non-negative as well. So far, no physical meaning is associated with these constants. It is obvious, that this split allows us to write the energy density (29) as

$$\rho = (1 + \kappa) \sqrt{\frac{A_2}{1 + \kappa} + \frac{B_2}{a^6}} . \quad (31)$$

This means, ρ can be regarded as consisting of two components, ρ_1 and ρ_2 ,

$$\rho = \rho_1 + \rho_2 , \quad (32)$$

with

$$\rho_1 = \sqrt{\frac{A_2 \kappa^2}{1 + \kappa} + \frac{B_2 \kappa^2}{a^6}} \quad (33)$$

and

$$\rho_2 = \sqrt{\frac{A_2}{1 + \kappa} + \frac{B_2}{a^6}} , \quad (34)$$

where $\rho_1 = \kappa \rho_2$ is valid, i.e., the ratio $\rho_1/\rho_2 = \kappa$ is constant. Obviously, both these expressions (33) and (34) describe energy densities for Chaplygin gases again. The corresponding (effective) equations of state are (the notations are chosen in agreement with the formalism of subsection 2.1)

$$p_1^* = -\frac{A_2 \kappa^2}{(1 + \kappa) \rho_1} = -\frac{A_2 \kappa}{(1 + \kappa) \rho_2} , \quad (35)$$

and

$$p_2^* = -\frac{A_2}{(1 + \kappa) \rho_2} . \quad (36)$$

This implies $p_1^* = \kappa p_2^*$, i.e., the effective pressures of both components differ by the same constant which also characterizes the ratio of both energy densities. It is straightforward to check that

$$p_1^* + p_2^* = -\frac{A_2}{\rho_2} = -\frac{A}{\rho} = p . \quad (37)$$

It is further convenient to introduce a quantity

$$p_2 \equiv -\frac{A_2}{\rho_2} . \quad (38)$$

This quantity differs from p_2^* by

$$p_2^* - p_2 = \frac{\kappa A_2}{(1 + \kappa) \rho_2} = -p_1^* . \quad (39)$$

Defining also a quantity Π by

$$\Pi \equiv p_1^* = -\frac{\kappa}{1 + \kappa} \frac{A_2}{\rho_2} = -\frac{\kappa}{1 + \kappa} \frac{A}{\rho} , \quad (40)$$

one checks by direct calculation that the following relations are valid:

$$\dot{\rho}_1 + 3H\rho_1 = -3H\Pi \quad \Leftrightarrow \quad \dot{\rho}_1 + 3H(\rho_1 + p_1^*) = 0 , \quad (41)$$

and

$$\dot{\rho}_2 + 3H(\rho_2 + p_2) = 3H\Pi \quad \Leftrightarrow \quad \dot{\rho}_2 + 3H(\rho_2 + p_2^*) = 0 . \quad (42)$$

Adding up these balances, consistently reproduces the total energy conservation

$$\dot{\rho} + 3H(\rho + p) = 0 . \quad (43)$$

At this point it becomes clear what we have obtained by the formal manipulations in Eq.(30). The set of equations (41) - (43) fits exactly the two-component description (6) - (8). It describes a two-component system in which the components interact with each other. The role of the interaction is to keep the ratio κ of the energy densities of both components constant. In other words, *any given Chaplygin gas can be thought as being composed of another Chaplygin gas which is in interaction with a pressureless fluid such that the energy density ratio of both components is fixed*. Apparently, this admits a further decomposition of the resulting Chaplygin gas component 2 in which now A_2 plays the role that A played so far, into still another Chaplygin gas interacting with matter in a similar way etc. It will turn out, however, that this formal possibility is not relevant for our approach.

3.2. Chaplygin gas interacting with matter

For our further considerations it is essential that the entire reasoning of the previous subsection 3.1 may be reversed. In such a case the starting point is a mixture of a Chaplygin gas and a pressureless fluid which are in interaction with each other according to

$$\dot{\rho}_1 + 3H\rho_1 = -3H\tilde{\Pi} , \quad (44)$$

$$\dot{\rho}_2 + 3H(\rho_2 + p_2) = 3H\tilde{\Pi} , \quad (45)$$

with

$$p_2 = -\frac{A_2}{\rho_2} = p \quad (46)$$

and an (initially unspecified) interaction quantity $\tilde{\Pi}$. This is a special case of (6), (7). For the time dependence of the (at this moment still arbitrary) ratio $\kappa \equiv \rho_1/\rho_2$ we have

$$\dot{\kappa} = \kappa \left[\frac{\dot{\rho}_1}{\rho_1} - \frac{\dot{\rho}_2}{\rho_2} \right] = 3H\kappa \left[\frac{p_2}{\rho_2} - \frac{\rho}{\rho_1\rho_2}\tilde{\Pi} \right] . \quad (47)$$

The stationarity requirement for κ then implies

$$\dot{\kappa} = 0 \quad \Leftrightarrow \quad \tilde{\Pi} = \frac{\rho_1}{\rho} p_2 . \quad (48)$$

With (46) and $\rho_1/\rho = \kappa/(1 + \kappa)$ one recovers relation (40) if $\tilde{\Pi}$ is identified with Π , i.e.,

$$\tilde{\Pi} = \Pi = -\frac{\kappa}{1 + \kappa} \frac{A_2}{\rho_2} = -\frac{\kappa}{1 + \kappa} \frac{A}{\rho} = -\kappa \frac{A_2}{\rho} . \quad (49)$$

Consequently, the energy densities of the components are again given by (33) and (34) and add up to (31) which, by redefining the constants according to (30), results in the energy density (29).

To summarize, we may either follow the line of the previous subsection 3.1 and find that one may attribute a specific two-component interpretation to any given Chaplygin gas or we may, alternatively, assert that a system consisting of a Chaplygin gas and a pressureless fluid which are in interaction with each other according to (44), (45) and (48), is equivalent to another Chaplygin gas with the energy density (29).

The components have sound velocities which are different from each other and different from the overall sound velocity. For the latter we have

$$\frac{\dot{p}}{\dot{\rho}} = -\frac{p}{\rho} = \frac{A}{\rho^2} = \frac{A}{A + \frac{B}{a^6}}. \quad (50)$$

The sound velocities of the components are $\dot{p}_1/\dot{\rho}_1 = 0$ and

$$\frac{\dot{p}_2}{\dot{\rho}_2} = -\frac{p_2}{\rho_2} = \frac{A_2}{\rho_2^2} = \frac{A_2}{\frac{A_2}{1+\kappa} + \frac{B_2}{a^6}} = (1 + \kappa) \frac{\dot{p}}{\dot{\rho}}. \quad (51)$$

Since for large times we have $\dot{p}/\dot{\rho} \rightarrow 1$, the quantity $\dot{p}_2/\dot{\rho}_2$ may be larger than unity which at the first glance seems to imply a superluminal sound propagation. However, $\dot{p}_2/\dot{\rho}_2$ is a formal quantity only, which does not describe any propagation phenomenon. The adiabatic sound speed (50) may also be split with respect to the effective pressures according to (10) with

$$\frac{\dot{p}_1^*}{\dot{\rho}_1} = \frac{\dot{p}}{\dot{\rho}} \quad \text{and} \quad \frac{\dot{p}_2^*}{\dot{\rho}_2} = \frac{\dot{p}}{\dot{\rho}}. \quad (52)$$

These effective sound velocities of the components coincide with the total sound velocity.

4. Non-adiabatic pressure perturbations

4.1. Perturbation dynamics

4.1.1. Effective sound speed

The Chaplygin in its two-component interpretation of the previous section belongs to a class of models with

$$\dot{\kappa} = 0 \quad \text{and} \quad \rho_1 + p_1^* = \kappa (\rho_2 + p_2^*). \quad (53)$$

Generally, the components may have (not necessarily constant) equations of state

$$p_1 = w_1 \rho_1 \quad \text{and} \quad p_2 = w_2 \rho_2. \quad (54)$$

For the total equation of state parameter w it follows that

$$w = \frac{w_2 + \kappa w_1}{1 + \kappa} \quad \text{and} \quad p = w \rho. \quad (55)$$

For our special Chaplygin gas case these quantities are

$$w_1 = 0, \quad w_2 = -\frac{A_2}{\rho_2^2}, \quad w = -\frac{A}{\rho^2}. \quad (56)$$

Under the condition (53) the difference of the individual sound speeds is

$$\frac{\dot{p}_2}{\dot{\rho}_2} - \frac{\dot{p}_1}{\dot{\rho}_1} = \frac{1 + \kappa}{\kappa} \frac{\dot{\Pi}}{\dot{\rho}_2} . \quad (57)$$

Introducing now matter perturbations in terms of the quantities defined in subsection 2.2 and allowing the density ratio κ to fluctuate, the difference between the fractional energy density perturbations which describes entropy perturbations becomes directly proportional to $\hat{\kappa}$:

$$D_2 - D_1 = 3(\zeta_2 - \zeta_1) = -\frac{1}{1 + w} \frac{\hat{\kappa}}{\kappa} . \quad (58)$$

This demonstrates, that a perturbation of the density ratio is essential for entropy perturbations to occur. Of course, for a “true” constant κ there are no non-adiabatic contributions and the two-component description does not give anything new compared to the original one-component picture. It should be mentioned that, since $\dot{\kappa} = 0$, the quantity $\hat{\kappa}$ is gauge-invariant (as is the combination on the left-hand side). Since the background ratio κ also determines the relation between the equation of state parameters A and A_2 in (30) a further specification may be performed. We shall assume from now on the parameter A_2 to be a true constant, i.e., $\hat{A}_2 = 0$. This choice implies

$$\frac{\hat{A}}{A} = \frac{\hat{\kappa}}{1 + \kappa} . \quad (59)$$

Within this model a fluctuation of the density ratio is equivalent to a fluctuation of the equation of state parameter A . Any fluctuation of κ or A generates a non-adiabatic pressure fluctuation

$$P - \frac{\dot{p}}{\dot{\rho}} D = \frac{w}{1 + w} \frac{\hat{\kappa}}{1 + \kappa} = \frac{w}{1 + w} \frac{\hat{A}}{A} . \quad (60)$$

We conclude that perturbing the equation of state parameter A represents a way to introduce non-adiabatic pressure perturbations in a Chaplygin gas. It should be mentioned that this type of perturbation can be obtained in a much simpler way by admitting fluctuations of A in the equation of state (28) itself. It is the most obvious choice which strictly speaking does not depend on the assumption of a specific internal structure. Nevertheless, we believe the two fluid context chosen here to be superior since it is connected with a model for the origin of the fluctuations of the equation of state parameter. Moreover, it is expedient to recall that in an underlying string theoretical formalism the parameter A is connected with the interaction strength of d-branes [5]. Hence, a fluctuating A corresponds to fluctuating interactions in string theory. An alternative, perhaps more transparent way of understanding the meaning of the fluctuation of the parameter A is to consider the Born-Infeld action that leads to the Chaplygin gas. The Lagrangian density in this case takes the form [35, 36]

$$L = \sqrt{-g} V(\phi) \sqrt{-\det[g_{\mu\nu} + \phi_{;\mu} \phi_{;\nu}]} , \quad (61)$$

where $V(\phi)$ is a potential term. This Lagrangian density leads to the Chaplygin gas equation of state for [8, 10]

$$V(\phi) = \sqrt{A} . \quad (62)$$

For $\phi = \phi_0 + \hat{\phi}$, the fluctuation in A is given by

$$\frac{\hat{A}}{A} = 2 \left(\frac{V'}{V} \right)_{\phi=\phi_0} \hat{\phi} \quad , \quad (63)$$

where the prime denotes the derivative with respect to ϕ . Consequently, fluctuations of A are allowed if the scalar field ϕ is not in the minimum of the potential. It has been shown that there are configurations for which a sufficiently flat potential (equivalent to an almost constant A) admits accelerated expansion [36].

For vanishing fluctuations of κ and A in (60) we recover the adiabatic case. Given that $w < 0$, any $\hat{A} > 0$ will reduce the adiabatic pressure perturbations. For definite statements further assumptions about $\hat{A}$ (or $\hat{\kappa}$) are necessary since otherwise the problem is undetermined. A simple case which will be used below is $\hat{A}/A = \text{constant}$. The structure of (60) also motivates a choice $\hat{A}/A = \mu(1+w)D$, for which

$$P = c_{eff}^2 D \quad , \quad c_{eff}^2 \equiv \frac{\dot{p}}{\dot{\rho}} (1 - \mu) \quad . \quad (64)$$

Any μ in the range $0 < \mu \leq 1$ leads to an effective sound speed square c_{eff}^2 which is reduced compared to the adiabatic value $\dot{p}/\dot{\rho} = -w$. It is interesting to note that the relation

$$\hat{p} = (1 - \mu) \frac{\dot{p}}{\dot{\rho}} \hat{\rho} = - (1 - \mu) \frac{p}{\rho} \hat{\rho} \quad ,$$

which is a different way of writing (64), coincides with that of a generalized Chaplygin gas with an equation of state $p = -C/\rho^{(1-\mu)}$ with (a true) constant C . For a generalized Chaplygin gas one would, however, also have $\dot{p} = - (1 - \mu) \frac{p}{\rho} \dot{\rho}$ and hence adiabatic perturbations only. In a sense, our strategy to consider perturbations $\hat{A} \neq 0$ while $\dot{A} = 0$, implies that the medium behaves as a Chaplygin gas in the background and shares features of a generalized Chaplygin gas on the perturbative level.

At this point also the role of the assumption $\hat{A}_2 = 0$ that precedes eq. (59) becomes clear. We mentioned at the end of subsection 3.1 that the decomposition of a given Chaplygin gas into another Chaplygin gas interacting with matter may be repeated again and again. In a subsequent step A_2 would play the role that A played so far. Hence, the assumption $\hat{A}_2 = 0$ ensures that no further non-adiabatic contributions will appear.

As already mentioned, a different two-fluid model for a (generalized) Chaplygin gas has been developed in reference [10], where this gas has been decomposed into a pressureless fluid and a time dependent cosmological “constant” with interactions between them. These authors obtained a modification of the perturbation dynamics through the interaction as well. The main difference to our approach is that in [10] the (generalized) Chaplygin gas was assumed to decompose into two components with constant equation of state parameters but variable ratio of the energy densities. Our splitting neither assumes the existence of a pressureless component from the outset nor does it restrict the equation of state parameters to have constant values. The split (32) - (34) which leads to (40) - (43) [with (28) and (29)] does not imply any a priori

requirements on the equation of state parameters of the components. The equation of state parameters in (56) are not imposed but emerge in our splitting procedure. No phantomlike contribution does occur in our setting which, moreover, is characterized by a different sign of the energy flow between the components. This shows that the claim of the authors of [10] to present a *unique* decomposition of the (generalized) Chaplygin gas into DM and DE is not generally true. It is only valid under the *additional* assumptions $p_{dm} = 0$ and $w_X = \text{const}$. Due to the circumstance that they deal with a Λ like fluid as DE component, perturbations of the latter do not occur in their analysis which seems to exclude perturbations of the parameter A .

4.1.2. Fluctuating decay rate

There exists still another interpretation for the appearance of non-adiabatic perturbations. The sign of the effective pressure Π in (49) indicates that within the two-component interpretation of the Chaplygin gas there is a transfer of energy from component 2 to component 1. This transfer may be described in terms of a decay rate Γ , defined by

$$3H\Pi \equiv -\Gamma\rho_2, \quad (65)$$

with the help of which the balance (45) may be written as

$$\dot{\rho}_2 + 3H(1 + w_2)\rho_2 = -\Gamma\rho_2. \quad (66)$$

From (49) we obtain $\Pi\rho = -\kappa A_2 \Rightarrow (\Pi\rho)^\cdot = 0$. On the other hand, (65) implies $\Pi\rho = -(\Gamma/3H)\rho_2\rho$, i.e., we have

$$\Pi\rho = -\frac{\Gamma}{3H} \frac{\rho^2}{1 + \kappa}. \quad (67)$$

Since in a spatially flat universe $\rho \propto H^2$ by Friedmann's equation, it follows that

$$(\Pi\rho)^\cdot = 0 \quad \Rightarrow \quad (\Gamma H^3)^\cdot = 0 \quad \Rightarrow \quad \Gamma \propto H^{-3}. \quad (68)$$

Consequently, a stationary ratio of the energy densities is guaranteed for a decay rate which scales with the inverse third power of the Hubble rate. *In the background, a Chaplygin gas with equation of state $p = -A/\rho$ can always be modelled as a mixture of a Chaplygin gas with equation of state $p_2 = -A_2/\rho_2$ and matter with $p_1 \ll \rho_1$, in which the Chaplygin gas component decays into the matter component at a rate $\Gamma \propto H^{-3}$.* Fluctuations of this decay rate are accompanied by fluctuations $\hat{\kappa}$ of the energy density ratio. Since $\hat{A}_2 = 0$ was assumed, we have

$$(\Pi\rho)^\cdot = -\hat{\kappa}A_2 \quad \Rightarrow \quad \frac{(\Pi\rho)^\cdot}{\Pi\rho} = \frac{\hat{\kappa}}{\kappa}. \quad (69)$$

An alternative expression, derived from (67), is

$$\frac{(\Pi\rho)^\cdot}{\Pi\rho} = \left[\frac{\hat{\Gamma}}{\Gamma} - \frac{\hat{H}}{H} + \frac{2\hat{\rho}}{\rho} - \frac{\hat{\kappa}}{1 + \kappa} \right]. \quad (70)$$

(Here $\hat{H}$ is defined as $\hat{\Theta}/3$, where $\Theta \equiv u_{;a}^a$ is the fluid expansion). On large perturbation scales for which spatial gradients may be neglected, the first order G_0^0 field equation implies

$$\frac{\hat{\rho}}{\rho} = 2 \frac{\hat{H}}{H} , \quad (71)$$

i.e.,

$$\frac{(\Pi\rho)^\wedge}{\Pi\rho} = \frac{(\Gamma H^3)^\wedge}{\Gamma H^3} - \frac{\hat{\kappa}}{1+\kappa} . \quad (72)$$

Combination of the two expressions (69) and (72) for $(\Pi\rho)^\wedge/(\Pi\rho)$ yields

$$\frac{(\Gamma H^3)^\wedge}{\Gamma H^3} = \frac{2+\kappa}{1+\kappa} \frac{\hat{\kappa}}{\kappa} . \quad (73)$$

Large scale fluctuations about $\Gamma \propto H^{-3}$ correspond to fluctuations of κ which, in turn, are equivalent to fluctuations of A according to (59).

4.1.3. Dynamics on large perturbation scales

Using the expression (60) in (25) we may now consider the large scale perturbation dynamics. If we neglect spatial gradients, the equation for ζ reduces to

$$\dot{\zeta} = -H \frac{w}{1+w} \frac{\hat{\kappa}}{1+\kappa} = -H \frac{w}{1+w} \frac{\hat{A}}{A} . \quad (74)$$

For a qualitative discussion of the impact of a non-vanishing fluctuation of the parameter A on the perturbation dynamics we assume $\hat{A}/A = \text{constant}$ as the simplest choice. With the Chaplygin gas equation of state

$$w = -\frac{Aa^6}{Aa^6 + B} \quad \Rightarrow \quad 1+w = \frac{B}{Aa^6 + B} \quad (75)$$

we obtain

$$\zeta = \zeta_i + \frac{1}{6} \frac{Aa^6}{B} \frac{\hat{A}}{A} . \quad (76)$$

The adiabatic case $\zeta = \zeta_i$ is, of course recovered for $\hat{A} = 0$. For $B \gg Aa^6$, i.e., for small a with a negligible pressure we have $\zeta \approx \zeta_i$. In the opposite limit $Aa^6 \gg B$ the non-adiabatic contribution can be substantial and may dominate the entire dynamics.

4.2. The gravitational potential

A time dependent equation of state and/or a varying ζ due to non-adiabatic perturbations generate a time dependence in the gravitational potential which, in turn, gives rise to the ISW. With the gauge-invariant definitions

$$\psi^x \equiv \psi + H\chi , \quad \psi^v \equiv \psi - Hv , \quad \phi^x \equiv \phi - \dot{\chi} , \quad (77)$$

where ψ^v is the comoving curvature perturbation, the relations (cf (4.92) in [37])

$$\zeta + \psi^v = \frac{2}{9(1+w)} \frac{\Delta\psi^x}{H^2 a^2} , \quad \phi^x = \psi^x \quad (78)$$

and (cf (4.169) in [37])

$$\dot{\psi}^\chi + \frac{3}{2}H \left(\frac{5}{3} + w \right) \psi^\chi = \frac{3}{2}(1+w)H\psi^v, \quad (79)$$

are generally valid. (The upper indices χ and v are kept to indicate the gauge-invariant character of the introduced quantities). From the definitions (23) for ζ and (77) for ψ^v it is obvious, that the combination $\zeta + \psi^v$ on the left-hand side of the first relation (78) describes comoving, fractional energy density perturbations. On large perturbation scales we may neglect the gradient term in (78) and ψ^v in (79) may be replaced by ζ :

$$\psi^v \approx -\zeta \quad \Rightarrow \quad \dot{\psi}^\chi + \frac{3}{2}H \left(\frac{5}{3} + w \right) \psi^\chi = -\frac{3}{2}(1+w)H\zeta. \quad (80)$$

A non-vanishing time dependence of the gravitational potential ψ^χ generates the ISW. The impact of the dark energy sound speed on the ISW has recently been discussed for quintessential models [17, 18, 19]. For (75) and (76) the equation for ψ^χ in (80) has the solution

$$\begin{aligned} \psi^\chi = & -\frac{3}{2}\zeta_i \frac{B}{a} E^{-1/4} \int \frac{da}{Aa^6 + B} E^{1/4} \\ & - \frac{1}{4} \frac{\hat{A}}{A} \frac{1}{a} E^{-1/4} \int da \frac{Aa^6}{Aa^6 + B} E^{1/4}, \end{aligned} \quad (81)$$

where

$$E \equiv \frac{Aa_i^6 + B}{Aa^6 + B} \left(\frac{a}{a_i} \right)^6. \quad (82)$$

The subscript i denotes some reference epoch with $a_i \ll a$ in the era $B \gg Aa_i^6$. The first term in (81) is an adiabatic contribution. The fluctuations in A give rise to a second, non-adiabatic gravitational potential term. For a qualitative discussion it is useful to consider the limit of very large values of the scale factor, i.e., for $Aa^6 \gg B$, where the background equation of state is close to that corresponding to a cosmological constant. In this realm the solution (81) is approximately

$$\psi^\chi \approx -\frac{3}{10}\zeta_i \frac{a_i}{a} \frac{B}{Aa_i^6} - \frac{1}{4} \frac{\hat{A}}{A}. \quad (83)$$

The first term decays with a^{-1} as a result of the time dependence of the equation of state. However, due to the non-adiabatic pressure perturbation there exists a constant second term which sets a threshold below which ψ^χ does not further decay. For sufficiently large values of the scale factor the first term becomes negligible and the gravitational potential approaches a constant value.

It is instructive to compare the potential (83) with the corresponding result of the Λ CDM model. With

$$\rho_{\Lambda\text{CDM}} = \rho_M + \rho_\Lambda, \quad \rho_M = \rho_{M_i} \left(\frac{a_i}{a} \right)^3, \quad \rho_\Lambda = \text{const}, \quad (84)$$

where the subindex i now denotes an initial value with $\rho_{M_i} \gg \rho_\Lambda$ and

$$p_M \ll \rho_M, \quad p_\Lambda = -\rho_\Lambda, \quad \Rightarrow \quad w_{\Lambda\text{CDM}} = -\frac{1}{1 + \frac{\rho_{M_i}}{\rho_\Lambda} \left(\frac{a_i}{a} \right)^3}, \quad (85)$$

we have $\zeta = \zeta_i$, i.e., ζ is a conserved quantity. The gravitational potential in the long time limit $a \gg a_i$ becomes

$$\psi_{\Lambda CDM}^x \approx -\frac{3}{4}\zeta_i \frac{\rho_{M_i}}{\rho_\Lambda} \frac{a_i}{a}. \quad (86)$$

In this case there does not exist a threshold value, $\psi_{\Lambda CDM}^x$ simply decays. Since the change in the gravitational potential affects the anisotropy spectrum at the largest scales one expects a discrimination between models with different values of $\hat{A}/A$.

5. Conclusions

We have demonstrated that any Chaplygin gas in a homogeneous and isotropic Universe can be regarded as being composed of another Chaplygin gas which is in interaction with a pressureless fluid such that the background ratio of the energy densities of both components remains constant. This two-component interpretation allowed us to establish a simple model for non-adiabatic pressure perturbations as the result of a specific internal structure of the substratum. Within the one-component picture this internal structure manifests itself as a spatial fluctuation of the parameter A in the Chaplygin gas equation of state $p = -A/\rho$. Such fluctuations, which can be traced back to fluctuations of a tachyon field potential, modify the adiabatic sound speed of the medium which may shed new light on the status of Chaplygin gas models of the cosmic substratum. For the special case of a constant fractional perturbation $\hat{A}/A$ on large scales we calculated the deviation of the quantity ζ , which describes curvature perturbations on constant density hypersurfaces, from its (constant) adiabatic value. Furthermore, fluctuations $\hat{A}$ give rise to an additional contribution to the gravitational potential which we have determined. In the long time limit when the Chaplygin gas equation of state is close to that corresponding to a cosmological constant, this contribution sets a threshold value below which the gravitational potential does not decay. This is different from both the adiabatic Chaplygin gas and the Λ CDM model with purely decaying potentials and should be a potential tool to discriminate between different dark energy models via different ISW contributions to the lowest multipoles in the anisotropy spectrum of the cosmic microwave background. A more quantitative analysis of this feature will be given in a subsequent paper.

Acknowledgements

This work was partially supported by the Deutsche Forschungsgemeinschaft. Discussions with Jérôme Martin and Nelson Pinto-Neto are gratefully acknowledged. JCF thanks the Institut für Theoretische Physik, Universität zu Köln for hospitality during part of this work. WZ is grateful for hospitality by the Departamento de Física Universidade Federal do Espírito Santo.

- [1] Riess, A.G. et al., *Astron. J.* 116, 1009 (1998); Perlmutter, S. et al., *Astrophys. J.* 517, 565 (1999); Riess, A.G. et al., *Astrophys. J.* 607, 665 (2004).
- [2] Peebles, P.J.E and Ratra, B., *Rev. Mod. Phys.* 75, 559 (2002); Sahni, V., Preprint astro-ph/00403324.
- [3] Amendola, L., *Phys. Rev. D* 62, 043511 (2000); Amendola, L. and Tocchini-Valentini, *Phys. Rev. D* 64, 043509 (2001); Zimdahl, W., Pavón, D. and Chimento, L.P., *Phys. Lett. B* 521, 133 (2001); Zimdahl, W. and Pavón, D., *Gen. Rel. Grav.* 35, 413 (2003); Chimento, L.P., Jakubi, A.S., Pavón, D. and Zimdahl, W., *Phys. Rev. D* 67, 083513 (2003); Pavón, D., Sen, S., and Zimdahl, W., *JCAP* 05, 009 (2004).
- [4] Chaplygin, S., *Sci. Mem. Moscow Univ. Math. Phys.* 21, 1 (1904).
- [5] Jackiw, R., *A Particle Field Theorist's Lectures on Supersymmetric, Non-Abelian Fluid Mechanics and d-Branes*, Preprint physics/0010042.
- [6] Kamenshchik, A. Yu, Moschella, U. and Pasquier, V., *Phys. Lett. B* 511, 265 (2001).
- [7] Fabris, J.C., Gonçalves, S.V.B. and de Souza, P.E., *Gen. Rel. Grav.* 34, 53 (2002).
- [8] Bento, M.C., Bertolami, O. and Sen, A.A., *Phys. Rev. D* 66, 043507 (2002).
- [9] Carturan, D. and Finelli, F., *Phys. Rev. D* 68, 103501 (2003).
- [10] Bento, M.C., Bertolami, O. and Sen, A.A., *Phys. Rev. D* 70, 083519 (2004).
- [11] Sandvik, H.B., Tegmark, M., Zaldariaga, M. and Waga, I., *Phys. Rev. D* 69, 123524 (2004).
- [12] Amendola, L., Finelli, F., Burigana, C. and Carturan, D., *JCAP* 0307, 005 (2003).
- [13] Bean, R. and Doré, O., *Phys. Rev. D* 68, 023515 (2003).
- [14] Bento, M.C., Bertolami, O. and Sen, A.A., *Phys. Lett. B* 575, 172 (2003).
- [15] Balakin, A.B., Pavón, D., Schwarz, D.J. and Zimdahl, W., *New Journal of Physics* 5, 85 (2003).
- [16] Reis, R.R.R., Waga, I., Calvao, M.O. and Joras, S.E., *Phys. Rev. D* 68, 061302 (2003).
- [17] Erickson, J.K., Caldwell, R.R., Steinhardt, P.J., Armendariz-Picon, C. and Mukhanov, V., *Phys. Rev. Lett.* 88, 1121301 (2002).
- [18] DeDeo, S., Caldwell, R.R., and Steinhardt, P.J., *Phys. Rev. D* 67, 103509 (2003).
- [19] Bean, R. and Doré, O., *Phys. Rev. D* 69, 083503 (2004).
- [20] Moroi, T. and Takahashi, T., *Phys. Rev. Lett.* 92, 091301 (2004).
- [21] Abramo, L.R., Finelli, F. and Pereira, T.S., *Phys. Rev. D* 70, 063517 (2004).
- [22] Gordon, Ch. and Hu, W., Preprint astro-ph/0406496.
- [23] Sachs, R.K. and Wolfe, A.M., *Astrophys.J.* 147, 73 (1967).
- [24] Nolta, M.R. et al., *Astrophys.J.* 608, 10 (2004).
- [25] Boughn, S.P. and Crittenden, R.G., Preprint astro-ph/0404470.
- [26] Vielva, P., Martinez-Gonzales, E. and Tucci, M., Preprint astro-ph/0408252.
- [27] Contaldi, C.R., Peloso, M., Kofman, L. and Linde, A., *JCAP* 07, 002 (2003).
- [28] Bastero-Gil, M., Freese, K. and Mersini-Houghton, L., *Phys. Rev. D* 68, 123514 (2003).
- [29] Tsujikawa, S., Maartens, R. and Brandenberger, R., *Phys. Lett. B* 574, 141 (2003).
- [30] Koivisto, T., Kurki-Suonio, H. and Ravndal, F., *Phys. Rev. D* 71, 064027 (2005).
- [31] Bardeen, J.M., Steinhardt, P.J. and Turner, M.S., *Phys. Rev. D* 28, 679 (1983).
- [32] Lyth, D.H., *Phys. Rev. D* 31, 1792 (1985).
- [33] García-Bellido, J. and Wands, D., *Phys. Rev. D* 53, 5437 (1996).
- [34] Wands, D., Malik, K.A., Lyth, D.H. and Liddle, A.R., *Phys. Rev. D* 62, 043527 (2000).
- [35] Sen, A., *JHEP* 9910, 008 (1999).
- [36] Copeland, E.J., Garousi, M.R., Sami, M. and Tsujikawa, S., *Phys. Rev. D* 71, 043003 (2005).
- [37] Liddle, A.R. and Lyth, D.H., *Cosmological Inflation and Large-Scale Structure* Cambridge University Press, Cambridge 2000.